


**PHYSICS**  
**Work, Energy, Power and Collisions**



1. At any instant of time  $t$ , the displacement of any particle is given by  $2t - 1$  (SI unit) under the influence of force of 5N. The value of instantaneous power is (in SI unit):  
(2024)  
(a) 5 (b) 7  
(c) 6 (d) 10
2. Two bodies A and B of same mass undergo completely inelastic one-dimensional collision. The body A moves with velocity  $v_1$  while body B is at rest before collision. The velocity of the system after collision is  $v_2$ . The ratio  $v_1 : v_2$  is.  
(2024)  
(a) 2 : 1 (b) 4 : 1  
(c) 1 : 4 (d) 1 : 2
3. The potential energy of a long spring when stretched by 2 cm is  $U$ . If the spring is stretched by 8 cm, potential energy stored in it will be:  
(2023)  
(a)  $16U$  (b)  $2U$   
(c)  $4U$  (d)  $8U$
4. A bullet from a gun is fired on a rectangular wooden block with velocity  $u$ . When bullet travels 24 cm through the block along its length horizontally, velocity of bullet become  $\frac{u}{3}$ . Then it further penetrates into the block in the same direction before coming to rest exactly at the other end of the block. The total length of the block is:  
(2023)  
(a) 30 cm (b) 27 cm  
(c) 24 cm (d) 28 cm
5. An electric lift with a maximum load of 2000 kg (lift + passengers) is moving up with a constant speed of  $1.5 \text{ ms}^{-1}$ . The frictional force opposing the motion is 3000 N. The minimum power delivered by the motor to the lift in watts is: ( $g = 10 \text{ ms}^{-2}$ )  
(2022)  
(a) 20000 (b) 34500  
(c) 23500 (d) 23000
6. Water falls from a height of 60 m at the rate of 15 kg/s to operate a turbine. The losses due to frictional force are 10% of the input energy. How much power is generated by the turbine? ( $g = 10 \text{ m/s}^2$ )  
(2021)  
(a) 8.1 kW (b) 12.3 kW  
(c) 7.0 kW (d) 10.2 kW
7. A point mass 'm' is moved in a vertical circle of radius 'r' with the help of a string. The velocity of the mass is  $\sqrt{7gr}$  at the lowest point. The tension in the string at the lowest point is  
(2020 Covid Re-NEET)  
(a) 7 mg (b) 8 mg  
(c) 1 mg (d) 6 mg
8. Body A of mass  $4m$  moving with speed  $u$  collides with another body B of mass  $2m$ , at rest. The collision is head on and elastic in nature. After the collision the fraction of energy lost by the colliding body A is :  
(2019)  
(a)  $\frac{1}{9}$   
(b)  $\frac{8}{9}$   
(c)  $\frac{4}{9}$   
(d)  $\frac{5}{9}$
9. A mass  $m$  is attached to a thin wire and whirled in a vertical circle. The wire is most likely to break when:  
(2019)  
(a) The mass is at the highest point  
(b) The wire is horizontal  
(c) The mass is at the lowest point  
(d) Inclined at an angle of  $60^\circ$  from vertical
10. A force  $F = 20 + 10y$  acts on a particle in  $y$  direction where  $F$  is in newton and  $y$  in metre. Work done by this force to move the particle from  $y = 0$  to  $y = 1$  m is  
(2019)  
(a) 30 J  
(b) 5 J  
(c) 25 J  
(d) 20 J

11. A body initially at rest and sliding along a frictionless track from a height  $h$  (as shown in the figure) just completes a vertical circle of diameter  $AB = D$ . The height  $h$  is equal to:



(2018)

- (a)  $\frac{7}{5}D$  (b)  $D$   
(c)  $\frac{3}{2}D$  (d)  $\frac{5}{4}D$

12. A spring of force constant  $k$  is cut into lengths of ratio  $1 : 2 : 3$ . They are connected in series and the new force constant is  $K'$ . Then they are connected in parallel and force constant is  $K''$ . Then  $K' : K''$  is:

(2017 - Delhi)

- (a)  $1 : 9$  (b)  $1 : 11$   
(c)  $1 : 14$  (d)  $1 : 6$

13. Consider a drop of rain water having mass  $1 \text{ g}$  falling from a height of  $1 \text{ km}$ . It hits the ground with a speed of  $50 \text{ m/s}$ . Take ' $g$ ' constant with a value  $10 \text{ m/s}^2$ . The work done by the (i) gravitational force and the (ii) resistive force of air is:

(2017 - Delhi)

- (a) (i)  $1.25 \text{ J}$  (ii)  $-8.25 \text{ J}$   
(b) (i)  $100 \text{ J}$  (ii)  $8.75 \text{ J}$   
(c) (i)  $10 \text{ J}$  (ii)  $-8.75 \text{ J}$   
(d) (i)  $-10 \text{ J}$  (ii)  $-8.25 \text{ J}$





14. A body initially at rest, breaks up into two pieces of masses  $2M$  and  $3M$  respectively, together having a total kinetic energy  $E$ . The piece of mass  $2M$ , after breaking up, has a kinetic energy.

(2017 - Gujarat)

- (a)  $\frac{2E}{5}$   
(b)  $\frac{E}{2}$   
(c)  $\frac{E}{5}$   
(d)  $\frac{3E}{5}$

15. A body is moving unidirectionally under the influence of a source of constant power supplying energy. Which one of the graph correctly shows the variation of displacement ( $s$ ) with time ( $t$ )?

(2017 - Gujarat)



16. A particle moves from a point  $(-2\hat{i} + 5\hat{j})$  to  $(4\hat{i} + 3\hat{k})$  when a force of  $(4\hat{i} + 3\hat{j}) \text{ N}$  is applied.

How much work has been done by the force? (2016 - II)

- (a)  $5 \text{ J}$  (b)  $2 \text{ J}$   
(c)  $8 \text{ J}$  (d)  $11 \text{ J}$

17. Two identical balls A and B having velocities of  $0.5 \text{ m/s}$  and  $-0.3 \text{ m/s}$  respectively collide elastically in one dimension. The velocities of B and A after the collision respectively will be:

(2016 - II)

- (a)  $-0.3 \text{ m/s}$  and  $0.5 \text{ m/s}$   
(b)  $0.3 \text{ m/s}$  and  $0.5 \text{ m/s}$   
(c)  $-0.5 \text{ m/s}$  and  $0.3 \text{ m/s}$   
(d)  $0.5 \text{ m/s}$  and  $-0.3 \text{ m/s}$

18. A bullet of mass 10 g moving horizontally with a velocity of  $400 \text{ ms}^{-1}$  strikes a wooden block of mass 2 kg which is suspended by a light inextensible string of length 5 m. As a result, the center of gravity of the block is found to rise a vertical distance of 10 cm. The speed of the bullet after it emerges out horizontally from the block will be:

(2016 - II)

- (a)  $120 \text{ ms}^{-1}$  (b)  $160 \text{ ms}^{-1}$   
(c)  $100 \text{ ms}^{-1}$  (d)  $80 \text{ ms}^{-1}$

19. A body of mass 1 kg begins to move under the action of a time dependent force  $F = (2t\hat{i} + 3t^2\hat{j}) \text{ N}$ , where  $\hat{i}$  and  $\hat{j}$  are unit vectors along x and y axis. What power will be developed by the force at the time t?

(2016 - I)

- (a)  $(2t^2 + 3t^2)W$  (b)  $(2t^2 + 4t^4)W$   
(c)  $(2t^3 + 4t^4)W$  (d)  $(2t^3 + 3t^5)W$

20. A piece of ice falls from a height h so that it melts completely. Only one-quarter of the heat produced is absorbed by the ice and all energy of ice gets converted into heat during its fall. The value of h is [Latent heat of ice is  $3.4 \times 10^5 \text{ J/kg}$  and  $g = 10 \text{ N/kg}$ ]:

(2016 - I)

- (a) 34 km (b) 544 km  
(c) 136 km (d) 68 km

21. What is the minimum velocity with which a body of mass m must enter a vertical loop of radius R so that it can complete the loop?

(2016 - I)

- (a)  $\sqrt{gR}$  (b)  $\sqrt{2gR}$   
(c)  $\sqrt{3gR}$  (d)  $\sqrt{5gR}$

22. A particle of mass 10 g moves along a circle of radius 6.4 cm with a constant tangential acceleration. What is the magnitude of this acceleration if the kinetic energy of the particle becomes equal to  $8 \times 10^{-4} \text{ J}$  by the end of the second revolution after the beginning of the motion?

(2016 - I)

- (a)  $0.1 \text{ m/s}^2$   
(b)  $0.15 \text{ m/s}^2$   
(c)  $0.18 \text{ m/s}^2$   
(d)  $0.2 \text{ m/s}^2$

23. A particle of mass m is driven by a machine that delivers a constant power k watts. If the particle starts from rest, the force on the particle at time t is:

(2015)

- (a)  $\sqrt{mk} t^{-\frac{1}{2}}$  (b)  $\sqrt{2mk} t^{-\frac{1}{2}}$   
(c)  $\frac{1}{2} \sqrt{mk} t^{-\frac{1}{2}}$  (d)  $\sqrt{\frac{mk}{2}} t^{-\frac{1}{2}}$

24. Two similar springs P and Q have spring constants  $K_P$  and  $K_Q$  such that  $K_P > K_Q$ . They stretched first by the same amount (case a), then by the same force (case b). The work done by the springs  $W_P$  and  $W_Q$  are related as in case (a) and case (b), respectively:

(2015)

- (a)  $W_P = W_Q$ ;  $W_P = W_Q$   
(b)  $W_P > W_Q$ ;  $W_Q > W_P$   
(c)  $W_P < W_Q$ ;  $W_Q < W_P$   
(d)  $W_P = W_Q$ ;  $W_P > W_Q$

25. A block of mass 10 kg moving in x direction with a constant speed of  $10 \text{ ms}^{-1}$ , is subjected to a retarding force  $F = -0.1x \text{ J/m}$  during its travel from  $x = 20 \text{ m}$  to  $30 \text{ m}$ . Its final K.E. will be:

(2015)

- (a) 450 J (b) 275 J  
(c) 250 J (d) 475 J

26. Two particles of masses  $m_1, m_2$  move with initial velocities  $u_1$  and  $u_2$ . On collision, one of the particles get excited to higher level, after absorbing energy  $\varepsilon$ . If final velocities of particles be  $v_1$  and  $v_2$ , then we must have:

(2015)

- (a)  $\frac{1}{2} m_1 u_1^2 + \frac{1}{2} m_2 u_2^2 = \frac{1}{2} m_1 v_1^2 + \frac{1}{2} m_2 v_2^2 - \varepsilon$   
(b)  $\frac{1}{2} m_1 u_1^2 + \frac{1}{2} m_2 u_2^2 - \varepsilon = \frac{1}{2} m_1 v_1^2 + \frac{1}{2} m_2 v_2^2$   
(c)  $\frac{1}{2} m_1^2 u_1^2 + \frac{1}{2} m_2^2 u_2^2 + \varepsilon = \frac{1}{2} m_1^2 v_1^2 + \frac{1}{2} m_2^2 v_2^2$   
(d)  $m_1^2 u_1 + m_2^2 u_2 - \varepsilon = \frac{1}{2} m_1^2 v_1^2 + \frac{1}{2} m_2^2 v_2^2$

27. The heart of a man pumps 5 litres of blood through the arteries per minute at a pressure of 150 mm of mercury. If the density of mercury be  $13.6 \times 10^3 \text{ kg/m}^3$  and  $g = 10 \text{ m/s}^2$ , then the power of heart in watt is:

(2015 Re)

- (a) 1.50  
(b) 1.70  
(c) 2.35  
(d) 3.0

- 28.** On a frictionless surface, a block of mass  $M$  moving at speed  $v$  collides elastically with another block of same mass  $M$  which is initially at rest. After collision the first block moves at an angle  $\theta$  to its initial direction and has a speed  $v/3$ . The second block's speed after the collision is: **(2015 Re)**
- (a)  $\frac{\sqrt{3}}{2}v$  (b)  $\frac{2\sqrt{2}}{3}v$   
 (c)  $\frac{3}{4}v$  (d)  $\frac{3}{\sqrt{2}}v$
- 29.** A ball is thrown vertically downwards from a height of 20 m with an initial velocity  $u_0$ . It collides with the ground, loses 50 percent of its energy in collision and rebounds to the same height. The initial velocity  $u_0$  is: (Take  $g = 10 \text{ ms}^{-2}$ ) **(2015 Re)**
- (a) 10 m/s (b) 14 m/s  
 (c) 20 m/s (d) 28 m/s
- 30.** Two particles A and B, move with constant motion in one dimension with velocities  $\vec{v}_1$  and  $\vec{v}_2$ . At the initial moment their position vectors are  $\vec{r}_1$  and  $\vec{r}_2$  respectively. The condition for particle A and B for their collision is: **(2015 Re)**
- (a)  $\vec{r}_1 - \vec{r}_2 = \vec{v}_1 - \vec{v}_2$   
 (b)  $\frac{\vec{r}_1 - \vec{r}_2}{|\vec{r}_1 - \vec{r}_2|} = \frac{\vec{v}_2 - \vec{v}_1}{|\vec{v}_2 - \vec{v}_1|}$   
 (c)  $\vec{r}_1 \cdot \vec{v}_1 = \vec{r}_2 \cdot \vec{v}_2$   
 (d)  $\vec{r}_1 \cdot \vec{v}_1 = \vec{r}_2 \cdot \vec{v}_2$
- 31.** If vectors  $\vec{A} = \cos\omega t\hat{i} + \sin\omega t\hat{j}$  and  $\vec{B} = \cos\frac{\omega t}{2}\hat{i} + \sin\frac{\omega t}{2}\hat{j}$  are functions of time, then the value of  $t$  at which they are orthogonal to each other are: **(2015 Re)**
- (a)  $t = 0$  (b)  $t = \frac{\pi}{4\omega}$   
 (c)  $t = \frac{\pi}{2\omega}$  (d)  $t = \frac{\pi}{\omega}$
- 32.** A body of mass  $(4m)$  is lying in x-y plane at rest. It suddenly explodes into three pieces. Two pieces, each of mass  $(m)$  move perpendicular to each other with equal speeds  $(v)$ . The total kinetic energy generated due to explosion is **(2014)**
- (a)  $mv^2$  (b)  $\frac{3}{2}mv^2$   
 (c)  $2mv^2$  (d)  $4mv^2$
- 33.** A uniform force of  $(3\hat{i} + \hat{j})$  newton acts on a particle of mass 2 kg. Hence the particle is displaced from position  $(2\hat{i} + \hat{k})$  metre to position  $(4\hat{i} + 3\hat{j} - \hat{k})$  metre. The work done by the force on the particle is: **(2013)**
- (a) 15 J (b) 9 J  
 (c) 6 J (d) 13 J
- 34.** The upper half of an inclined plane of inclination  $\theta$  is perfectly smooth while lower half is rough. A block starting from rest at the top of the plane will again come to rest at the bottom, if the coefficient of friction between the block and lower half of the plane is given by: **(2013)**
- (a)  $\mu = \tan\theta$  (b)  $\mu = \frac{1}{\tan\theta}$   
 (c)  $\mu = \frac{2}{\tan\theta}$  (d)  $\mu = 2\tan\theta$