



DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

Test Booklet Series



TEST BOOKLET

MEDICAL OFFICER
(ASSISTANT SURGEON)

T. B. C. : AS – 18

SI. No. 2549

Time Allowed: 3 Hours

Maximum Marks : 200

: INSTRUCTIONS TO CANDIDATES :

- 1. IMMEDIATELY AFTER THE COMMENCEMENT OF THE EXAMINATION, YOU SHOULD CHECK THAT THIS TEST BOOKLET DOES NOT HAVE ANY UNPRINTED OR TORN OR MISSING PAGES OR ITEMS ETC. IF SO, GET IT REPLACED BY A COMPLETE TEST BOOKLET OF THE SAME SERIES ISSUED TO YOU.
- 2. ENCODE CLEARLY THE TEST BOOKLET SERIES A, B, C OR D, AS THE CASE MAY BE, IN THE APPROPRIATE PLACE IN THE ANSWER SHEET USING BALL POINT PEN (BLUE OR BLACK).
- 3. You have to enter your Roll No. on the Test Booklet in the Box provided alongside. DO NOT write anything else on the Test Booklet.
- 4. YOU ARE REQUIRED TO FILL UP & DARKEN ROLL NO., TEST BOOKLET / QUESTION BOOKLET SERIES IN THE ANSWER SHEET AS WELL AS FILL UP TEST BOOKLET / QUESTION BOOKLET SERIES AND SERIAL NO. AND ANSWER SHEET SERIAL NO. IN THE ATTENDANCE SHEET CAREFULLY. WRONGLY FILLED UP ANSWER SHEETS ARE LIABLE FOR REJECTION AT THE RISK OF THE CANDIDATE.
- 5. This Test Booklet contains 200 items (questions). Each item (question) comprises four responses (answers). You have to select the correct response (answer) which you want to mark (darken) on the Answer Sheet. In case, you feel that there is more than one correct response (answer), you should mark (darken) the response (answer) which you consider the best. In any case, choose ONLY ONE response (answer) for each item (question).
- 6. You have to mark (darken) all your responses (answers) ONLY on the separate Answer Sheet provided by using BALL POINT PEN (BLUE OR BLACK). See instructions in the Answer Sheet.
- 7. All items (questions) carry equal marks. All items (questions) are compulsory. Your total marks will depend only on the number of correct responses (answers) marked by you in the Answer Sheet. There will be no negative markings for wrong answers.
- 8. Before you proceed to mark (darken) in the Answer Sheet the responses to various items (questions) in the Test Booklet, you have to fill in some particulars in the Answer Sheet as per the instructions sent to you with your **Admission Certificate**.
- 9. After you have completed filling in all your responses (answers) on the Answer Sheet and after conclusion of the examination, you should hand over to the Invigilator the Answer Sheet issued to you. You are allowed to take with you the candidate's copy / second page of the Answer Sheet along with the Test Booklet, after completion of the examination, for your reference.
- 10. Sheets for rough work are appended in the Test Booklet at the end.

DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

Test Prime

ALL EXAMS, ONE SUBSCRIPTION



70,000+ Mock Tests



600+ Exam Covered



Personalised Report Card



Previous Year Papers



Unlimited Re-Attempt



500% Refund

















ATTEMPT FREE MOCK NOW





- In transverse section through superior mediastinum which structure is NOT seen?
 - (A) Arch of aorta
 - (B) Superior vena cava
 - (C) Oesophagus
 - (D) Left atrium
- 2. Atherosclerotic occlusion of which of the following arteries would result in insufficient perfusion of the urinary bladder?
 - (A) External iliac
 - (B) Inferior epigastric
 - (C) Internal iliac
 - (D) Internal pudendal
- 3. The following statements are true for great saphenous vein except:
 - (A) Travels with the saphenous nerve
 - (B) Passes posterior to the medial malleolus
 - (C) Ascends along the medial aspect of leg and thigh
 - (D) Empties into the femoral vein
- If breast carcinoma invades the retromammary space and pectoral fascia, contraction of which muscle may cause the whole breast to move superiorly.
 - (A) Pectoralis major
 - (B) Pectoralis minor
 - (C) Serratus anterior
 - (D) 2nd to 6th intercostal muscles

- 5. Cauda equina syndrome is characterized by all EXCEPT:
 - (A) Saddle anaesthesia
 - (B) Dissociate anaesthesia
 - (C) Asymmetric areflexic paraplegia
 - (D) Bowel and bladder disturbances
- 6. Melatonin secretion would probably not be increased by :
 - (A) Stimulation of superior cervical ganglia
 - (B) Intravenous infusion of tryptophan
 - (C) Intravenous infusion of epinephrine
 - (D) Stimulation of the optic nerve
- 7. A young woman has puffy skin and a hoarse voice. Her plasma TSH concentration is low but increases markedly when she is given TRH. She probably has:
 - (A) Hypot<mark>hyroidism due to a thyroid</mark> tumor
 - (B) Hypothyroidism due to a primary abnormality in the thyroid gland
 - (C) Hypothyroidism due to a primary abnormality in the pituitary gland
 - (D) Hypothyroidism due to primary abnormality in the hypothalamus





- Glucagon increases glycogenolysis in liver cells but ACTH does not because:
 - (A) Cortisol increases the plasma glucose level
 - (B) Liver cells have an adenylyl cyclase different from that in adrenocortical cells
 - (C) ACTH can not enter the nucleus of liver cells
 - (D) Membrane of liver cells contain receptors different from those in adrenocortical cells
- 9. Drugs and toxins that increase the CAMP content of intestinal mucosa cause diarrhea because they:
 - (A) Increase K⁺ secretion into colon
 - (B) Inhibit K⁺ absorption in crypts of Lieberkuhn
 - (C) Increase Na⁺ absorption in small intestine
 - (D) Increase CI secretion into intestinal lumen
- 10. The work performed by left ventricle is substantially greater than that performed by right ventricle, because in the left ventricle;
 - (A) Wall is thicker
 - (B) Stroke volume is greater

- (C) Preload is greater
- (D) After load is greater
- 11. Proteins targeted for destruction in eukaryotes are covalently linked to:
 - (A) Clathrin
 - (B) Pepsin
 - (C) Laminin
 - (D) Ubiquitin
- 12. An example of a tumour suppressor gene is :
 - (A) Myc
 - (B) Fos
 - (C) Ras
 - (D) Rb
- 13. All is true about glutathione except :
 - (A) It is a tripeptide
 - (B) It converts hemoglobin to methemoglobin
 - (C) It conjugates xenobiotics
 - (D) It scavenges free radicals and superoxide ions
- 14. The amino acid carboxylated by vitamin K is:
 - (A) Aspartate
 - (B) Glutamate
 - ∴ (C) Histidine
 - (D) Proline

(3)





- 15. Which one of the following is a membrane bound enzyme that catalyzes the formation of cyclic AMP from ATP?
 - (A) Tyrosine Kinase
 - (B) Polymerase
 - (C) Adenylate cyclase
 - (D) ATP synthase
- 16. Which of the following is responsible for Alzheimer's disease?
 - (A) PrP_{SC} Protein
 - (B) PrP_C Protein
 - (C) Amyloid β
 - (D) Parkin
- 17. Action on Nitric oxide is mediated through:
 - (A) cGMP
 - (B) cAMP
 - ' (C) Ca⁺⁺
 - (D) Diacylglycerol
- 18. Which of the following laboratory tests will be useful in patients of primary hypothyroidism?
 - (A) Estimation of T3
 - (B) Estimation of T4
 - (C) Estimation of TBG
 - (D) Detection of auto-antibodies
- 19. The lipoprotein with fastest electrophoretic mobility and lowest triglyceride content is:
 - (A) VLDL
 - (B) IDL

- (C) LDL
- (D) HDL
- 20. Hemolytic anemia in patients taking anti-malarial drug primaquine can be attributable to the deficiency of:
 - (A) Glycogen phosphorylase
 - (B) Glyceraldehyde 3-phosphate dehydrogenase
 - (C) Glucose 6-phosphate dehydrogenase
 - (D) Glucose 6-phosphatase
- 21. Anticholinesterate agent with selective effect on neuromuscular junction is:
 - (A) Neostigmin
 - (B) Physostigmin
 - (C) Parathione
 - (D) None of the above
- 22. Toxicity by loading dose is least seen on:
 - (A) Liver
 - (B) Kidney
 - (C) Brain
 - (D) Muscle
- 23. Pinocytosis is typically absent in capillary endothelium of :
 - (A) Liver
 - (B) Kidney
 - (C) Brain
 - (D) Muscle

(4)

Contd.





- 24. Major role in regulation of coronary blood flow is played by:
 - (A) Acetylcholine
 - (B) Noradrenaline
 - (C) Adenosine
 - (D) All of the above
- 25. In designing long term drug dosing schedule, the most important concept is:
 - (A) Bioavailability
 - (B) Apparent volume of distribution
 - (C) Clearance
 - (D) · All are equally important
- 26. Celiprolol is:
 - (A) Both alfa and beta adrenoceptor blocker
 - (B) Selective beta-1 blocker
 - (C) Non-selective beta blocker
 - (D) Beta-1 blocker and Beta-2 agonist
- 27. Order of receptors stimulated by increasing Dopamine concentration is:
 - (A) Dopamine-1, beta-1, alfa-1
 - (B) Dopamine-1, alfa-1, beta-1
 - (C) Alfa-1, dopamine-1, beta-1
 - (D) Beta-1, dopamine-1, alfa-1
- 28. Paracetamol toxicity include:
 - (A) Metabolic acidosis-
 - (B) Prolonged bleeding time.

- (C) Both (A) and (B)
- (D) None of the above
- 29. Thiazide does not induce increased:
 - (A) Uric acid excretion
 - (B) Potassium excretion
 - (C) Chloride excretion
 - (D) Calcium reabsorption
- 30. Antiepileptic mechanism of phenytoin involves :
 - (A) Promoting open state of chloride channel
 - (B) Glutamate antagonism
 - (C) Delaying recovery of inactive sodium channel
 - (D) All of the above
- 31. In an adult patient bleeding time is observed to be prolonged [12 minutes] tests for which of the following needs to be immediately performed?
 - (A) Extrinsic pathway clotting defect
 - (B) Intrinsic pathway clotting defect
 - (C) Platelet function/count
 - (D) Fibrinogen activity/level
- 32. Lines of Zahn help pathologist in the identification / diagnosis of :
 -(A) ,Thrombus
 - (B) Layers in tissue Infarct
 - (C) Post-mortem clot
 - (D) Zones of Gangrene

(*)

(5)





- 33. Sections from Brain Specimen of an aged female showed evidences of fat embolism, she likely suffered from:
 - (A) Head injury
 - (B) Long bone fracture
 - (C) Drowning
 - (D) Fat necrosis of Pancreas
- 34. Heart-failure Cells in the tissue section are diagnostic of the following:
 - (A) CVC Lung
 - (B) CVC Heart
 - (C) CVC Liver
 - (D) CVC Brain
- 35. Haemophilia is linked with the Chromosome:
 - (A) Chromosome 3
 - (B) Chromosome 16
 - (C) Chromosome X
 - (D) Chromosome Y
- 36. Rosette formations are the characteristicstic feature of :
 - (A) Dysgerminoma
 - (B) Melanoma
 - (C) Small-cell lymphoma
 - (D) Retinoblastoma
- 37. Myasthenia gravis is commonly associated with:
 - (A) Parathyroid Adenoma
 - (B) Thyroid Carcinoma

- (C) Thymoma
- (D) Basophilic Adenoma
- 38. Findings in a patient having Hb-6 gm%, TLC 1200/cumm, Platelet count 60,000/cumm, MCV 12fl are suggestive of diagnosis of:
 - (A) Aplastic Anaemia
 - (B) Megaloblastic Anaemia
 - (C) PNH
 - (D) Myelofibrosis
- 39. Aschoff's nodules are the diagnostic features of the following disease:
 - (A) Sub-acute bacterial Endocarditis
 - (B) Rheumatic Carditis
 - (C) Libman Sacks Endocarditis
 - (D) Thrombotic Endocarditis
- 40. Patient of kidney disease is diagnosed to be presenting with Oliguria if urinary output is:
 - (A) Less than 20 ml/24 hours
 - (B) Less than 40 ml/24 hours
 - (C) Less than 400 ml/24 hours
 - (D) Less than 800 ml/24 hours
- 41. Prozone phenomenon is a feature in:
 - (A) Tularemia
 - (B) Legionnaire's disease
 - (C) Plague
 - (D) Brucellosis

'Confd.





- 42. Minimum Inhibitory Concentration (MIC) can be calculated by all of the following antibiotic sensitivity methods except:
 - (A) Epsilometer test
 - (B) Agar dilution method
 - (C) Kirby Bauer's disk diffusion method
 - (D) Broth dilution method
- 43. Suckling mice is used for isolation of:
 - (A) Coxsackie virus
 - (B) Herpes virus
 - (C) Pox virus
 - (D) Adenovirus
- 44. Both Intracytoplasmic and Intranuclear inclusions are seen in:
 - (A) Herpes virus
 - (B) Adeno virus
 - (C) Measles virus
 - (D) HIV virus
- 45. Varicella-zoster remains latent in:
 - (A) Lymphocytes 4
 - (B) Monocytes
 - (C) Trigeminal ganglion
 - (D) Plasma cells
- 46. Antigenic variation seen in which of the following?
 - (A) Influenza virus
 - (B) Hepatitis virus

- (C) Yellow fever virus
- (D) Leptospira
- 47. In Japanese encephalitis, pigs act as:
 - (A) Definitive host
 - (B) Amplifier host
 - (C) Intermediate host
 - (D) Any of the above
- 48. Hepatitis A virus can be inactivated by:
 - (A) Boiling over 120°C over 1 minute
 - (B) Ether
 - (C). Chloroform
 - (D) Iodine
- 49. Important feature of AIDS is:
 - (A) Follicular tonsillitis
 - (B) Lichen planus
 - (C) Oral candidiasis
 - (D) Mitotic lesions of the oral cavity
- 50. Malarial parasite which statement is false regarding communicability?
 - (A) The gametocytes appear in blood 4-6 days after a sexual phase in P. vivax.
 - (B) The gametocytes appear in blood 10-12 days after a sexual phase in P. falciparum.
 - (C) The number of gametocytes increases in blood with time.
 - (D) The number of gametocytes increases by 1000 times.

(7)





- 51. Which one of the following is not followed in India?
 - (A) Recording a dying declaration
 - (B) Recording a dying deposition
 - (C) Conducting a policy inquest
 - (D) Conducting a magistrate's inquest
- 52. Vesicles in ante-mortem burn are differentiated from post-mortem burn by the presence of :
 - (A) Chloride
 - (B) Albuminous fluid and chloride
 - (C) Plasma and WBC in fluid
 - (D) Air
- 53. Somatic death is said to occur when:
 - (A) All individual cells of the person die
 - (B) The person is dead clinically, although many of his individual cells may still be functional
 - (C) The heart stops beating permanently
 - (D) The person is on artificial life support system
- 54. Gang rape has been defined in India law as:
 - (A) Rape committed on a woman by a member of a criminal gang
 - (B) Rape committed by a member in order to induce the woman to join their gang

- (C) A rape by more than one in a group of persons acting in furtherance of their common intention
- (D) Rape followed by killing of the woman
- 55. Pus formation in a lacerated wound takes place within:
 - (A) 6 to 13 hours
 - (B) 12 to 24 hours
 - (C) 36 to 48 hours
 - (D) 7 to 10 days
- 56. The greater cornu of hyoid bone unites with the body:
 - (A) Between 10 to 20 years
 - (B) Between 20 to 30 years
 - (C) Between 30 to 40 years
 - (D) Between 40 to 60 years
- 57. Which of the following sign will indicate that the menstrual blood was ante-mortem?
 - (A) Alkaline
 - (B) Light pink
 - (C) Does not clot unless mixed with seminal fluid
 - (D) Shows endometrial and vaginal cells
- 58. 'Brush burn' is:
 - (A) An abrasion
 - (B) An electric burn
 - (C) A chemical burn
 - (D) None of the above





- 59. For viriological examination, the tissue is preserved usually in:
 - (A) 10% formalin
 - (B) 50% glycerine
 - (C) Rectified spirit
 - (D) Saturated solution of sodium chloride
- 60. In a suspected case of air embolism, search for embolism is to be done in:
 - (A) Circle of wills
 - (B) Meningeal vessels
 - (C) Abdominal aorta.
 - (D) Right ventricle of heart
- 61. There is narcotic abuse case with jet black tongue. The cause is:
 - (A) Cocaine poisoning
 - (B) Morphine poisoning
 - (C) Cannabis toxicity
 - (D) LSD poisoning
- 62. In Rokitansky technique of postmortem:
 - (A) Organs are removed one by one
 - (B) In-situ dissection is done
 - (C) Thoracic and cervical organs, abdominal organs and the organs of urogenital systems are removed as separate organ blocks
 - (D) Thoracic, cervical, abdominal and pelvic organs are removed as one large organ block

- 63. Post-mortem luminescence in a decaying body is due to:
 - (A) Entamoeba coli
 - (B) Armillaria mellea
 - (C) Clostridium welchii
 - (D) Formation of methane
- 64. To develop cyanosis, minimum percentage reduced haemoglobin should be more than:
 - (A) 11 gm%
 - (B) 3 gm%
 - (C) 5 gm%
 - (D) 7 gm%
- 65. Exposure and abandonment of child less than 12 years of age by his parents may lead to their prosecution under Section:
 - (A) 301 IPC
 - (B) 317 IPC
 - (C) 318 IPC
 - (D) 363 IPC
- 66. Which of the following is the most logical sequence?
 - (A) Impairment-Disease-Disability-Handicap
 - (B) Disease-Impairment-Disability-Handicap
 - (C) Disease-Impairment-Handicap-Disability
 - (D) Disease-Handicap-Impairment-Disability

WD - 1A/80 (9) (Turn over)





- 67. Screening test for detection of carcinoma of cervix is which level of prevention?
 - (A) Primordial
 - (B) Primary
 - (C) Secondary
 - (D) Tertiary
- 68. The distance from airport or seaport which has to be kept free from all types of mosquitoes is:
 - (A) 400 m
 - (B) 500 m
 - (C) 1 km
 - (D) 100 m
- 69. Number of air changes in a living room per hour should be at least:
 - (A) 2-3
 - (B) 3-4
 - (C) 4-5
 - (D) 5-6
- 70. Acculturation means:
 - (A) Culture contact
 - (B) Study of the various cultures
 - (C) Cultural history of health and disease
 - (D) None of the above
- 71. Argemone oil contamination of mustard oil can be detected by:
 - (A) Phosphatase test
 - (B) Nitric acid test

- (C) Coliform test
- (D) Methylene blue test
- 72. Ujjawala Scheme is for prevention of:
 - (A) Child abuse
 - (B) Child trafficking
 - (C) Child labour
 - (D) Child marriage
- 73. Kishori Shakti Yojana (KSY) is:
 - (A) Empowerment of females under Maternity Benefit Scheme
 - (B) Adolescent girl's scheme . under ICDS
 - (C) Free and compulsory education for girl child
 - (D) Child care home scheme for female juvenile delinquents
- 74. Maximum recommended number of students in a school class room is:
 - (A) 30
 - (B) 35
 - (C) 40
 - (D) 50
- 75. In India maximum maternal mortality is due to :
 - (A) Hemorrhage
 - (B) Anemia
 - (C) Abortion
 - (D) Sepsis





- 76. Disinfecting action of chlorine on water is mainly due to:
 - (A) Hydrogen chloride
 - (B) Hypochlorous acid
 - (C) Hypochlorite ions
 - (D) Hydrogen ions
- 77. The number of live births per 1000 women in the reproductive age group in a given year refers to:
 - (A) Total Fertility Rate
 - (B) Gross Reproduction Rate
 - (C) Net Reproduction Rate
 - (D) General Fertility Rate
- 78. All are true about Janani Shishu Suraksha Karyakram (JSSK), except:
 - (A) Free diet to mother during hospital stay
 - (B) Free delivery
 - (C) Free transport from home to hospital and back
 - (D) Free treatment of sick infants upto 1 year
- 79. According to Suraksha Clinic in National AIDS Control Program, person coming with lower abdominal pain, the color code of kit in treatment is:
 - (A) White
 - (B) Yellow
 - (C) Green
 - (D) Grey

- 80. Which of the following is didactic method of health communication?
 - (A) Group discussion
 - (B) Workshop
 - (C) Lecture
 - (D) Panel discussion
- 81. Followings are the modified sebaceous glands, EXCEPT:
 - (A) Bartholin's glands
 - (B) Mammary glands
 - (C) Meibomian glands in eyes
 - (D) Tyson's glands of prepuce
- 82. Which of the following is **NOT** a constituent of sebum?
 - (A) Cholesterol esters
 - (B) Glycogen
 - (C) Glycerides
 - (D) Wax esters
- 83. Herpes virus group cause following diseases, EXCEPT:
 - (A) Chicken pox
 - (B) Cytomegalovirus infection
 - (C) Infectious Mononucleosis
 - (D) Molluscum contagiosum
- 84. Candida albicans is associated with:
 - (A) Benign migratory glossitis
 - (B) Median rhomboid glossitis
 - (C) Fissured tongue
 - (D) Fordyces spots

(11)





- 85. Receptors for IgE are present on :
 - (A) Basophils
 - (B) Eosinophils
 - (C) Plasma cell
 - (D) Polymorphs
- 86. Patient with a metallic foreign body in eye, which investigation is contraindicated?
 - (A) MRI
 - (B) USG
 - (C) X-RAY
 - (D) CT SCAN
- 87. All of them emit radiation except:
 - (A) CT SCAN .
 - (B) X-RAYS
 - (C) Intravenous Pyelography
 - (D) Ultrasound
- 88. Most radiodense tissue in the body is:
 - (A) Fluid
 - (B) Brain
 - (C) Soft tissue
 - (D) Bone
- 89. The Magnetic field in a MRI is measured in:
 - (A) Hounsfield unit
 - (B) MHz
 - (C) Tesla
 - (D) None of the above
- 90. T1, T2 IMAGES are the features of which diagnostic modality?
 - (A) USG
 - (B) MRI
 - (C) CT SCAN
 - (D) Radiography

- 91. Choose the most effective mechanical device for cleaning concave interdental surfaces of teeth:
 - (A) Conventional tooth brushes
 - (B) Powered tooth brushes
 - (C) Triangular tooth picks
 - (D) Interdental brushes
- 92. Interproximal necrosis, ulceration, pain and bleeding are the significant criteria used in the clinical diagnosis of:
 - (A) Lichen Planus
 - (B) ANUG
 - (C) Gingival enlargement
 - (D) Chronic infectious periodontal disease
- 93. The predominant cell type in gingival (crevicular) fluid is the :
 - (A) Mast cell
 - (B) Plasma cell
 - (C) Macrophage
 - (D) PMN leukocyte
- 94. Important diagnostic tool for achieving accurate implant angulation:
 - (A) Diagnostic template
 - (B) Waxup
 - (C) Mounted diagnostic cast
 - (D) Surgical template:
- 95. Facebow is used to transfer:
 - (A) Axis orbital plane
 - (B) Frankfort horizon plane
 - (C) Camper's plane
 - (D) Occlusal plane





- 96. The antibody most specific for Systemic Lupus Erythematosus is:
 - (A) Anti RNA
 - (B) Anti SS-DNA
 - (C) Anti DS-DNA
 - (D) Anti Sm antibody
- 97. Raised intracranial tension is indicated by:
 - (A) Hypotension and bradycardia
 - (B) Hypotension and tachycardia
 - (C) Hypertension and bradycardia
 - (D) Hypertension and tachycardia
- 98. Patients with hereditary spherocytosis are more prone to develop:
 - (A) Malabsorption
 - (B) Recurrent UTI
 - (C) Malaria
 - (D) Gall stones
- 99. The adult hemoglobin consists of the following chains:
 - (A) $2\alpha + 2\beta$
 - (B) $2\alpha + 2\delta$
 - (C) $2\beta + 2\delta$
 - (D) $2\alpha + 2\gamma$
- 100. Mononeuritis multiplex is seen with:
 - (A) Leprosy
 - (B) Syphilis
 - (C) GB syndrome
 - (D) Vitamin B12 deficiency
- 101. A patient presents to the medical emergency with features of urinary retention, tachycardia and dry mouth

- after ingestion of an unidentified substance. The most likely etiology of poisoning in this patient is due to:
- (A) Quinine
- (B) Opioids
- (C) Organophosphorous compounds
- (D) Dhatura
- 102. Spike and dome pattern on an EEG is the characteristic of:
 - (A) Petit mal seizures
 - (B) Grand mal seizures
 - (C) Temporal lobe epilepsy
 - (D) Absence seizures
- 103. Keratoderma blenorrhagica is pathognomonic of:
 - (A) Behcet's disease
 - (B) Reiter's syndrome
 - (C) Lyme's disease
 - (D) Glucogonoma
- 104. Drug of choice for the treatment of Pneumocystis jiroveci pneumonia
 - is:
 - (A) Ceftriaxone
 - (B) Gentamicin
 - (C) Cotrimoxazole
 - (D) Azithromycin
- 105. Lactic acidosis is a recognized adverse effect of which of the following antiretroviral agents?
 - (A) Nevirapine
 - (B) Efavirenz
 - (C) Stayudine
 - (D) Lopinavir

(13)





- 106. Chloroma is associated with which of the following conditions?
 - (A) Dehydration
 - (B) Melanoma
 - (C) Myeloid leukemia
 - (D) Wilm's tumour
- 107. The volume of air taken into the lungs during normal breathing is called:
 - (A) Vital capacity
 - (B) Inspiratory reserve volume
 - (C) Tidal volume
 - (D) Maximum inspiratory capacity
- 108. The commonest site of a hypertensive intracranial bleed is:
 - (A) Thalamus
 - (B) Putamen
 - (C) Frontal lobe.
 - (D) Pons
- 109. Heberden's nodes are seen in :
 - (A) Acute rheumatic fever
 - (B) Rheumatoid arthritis
 - (C) Osteoarthritis
 - (D) Infective endocarditis
- 110. Mee's lines are seen in the nails in poisoning due to:
 - (A) Lead
 - (B) Arsenic
 - (C) Mercury
 - (D) Copper
- 111. The silhouette sign on a chest skiagram is seen in disease of the :
 - (A) Right lower lobe
 - (B) Lingual

- (C) Left upper lobe
- (D) Right middle lobe
- 112. The drug of choice for the treatment of cerebral malaria in a pregnant woman is:
 - (A) Chloroquin
 - (B) Mefloquin
 - (C) Quinine
 - (D) Pyrimethamine sulfadoxine
- 113. The presence of bronchial breath sounds and aegophony in a patient with pericardial effusion is called:
 - (A) Levine sign
 - (B) Kussmaul sign
 - (C) Ewart's sign
 - (D) Hill's sign
- 114. The most common causative organism of acute infective endocarditis is:
 - (A) Staphylococcus sp.
 - (B) Streptococcus viridans
 - (C) Enterococci
 - (D) Gram negative bacteria
- 115. Werniecke's encephalopathy is due to the deficiency of:
 - (A) Niacin
 - (B) Folate
 - (C) Vitamin B12
 - (D) Thiamine
- 116. Throughout the world the commonest cause of ill health is:
 - (A) Malignant disease
 - (B) Disease due to infection
 - (C) Malnutrition
 - (D) Degenerative arterial disease





117. Neonates are particularly susceptible to:

- (A) Scarlet fever
- (B) Measles
- (C) Whooping-cough
- (D) Rubella
- 118. The most common cause of acute circulatory failure (shock) is:
 - (A) Bacteraemia
 - (B) Massive pulmonary embolism
 - (C) Loss of fluid
 - (D) Myocardial infarction
- 119. During pregnancy significant cardiac disease is suggested by :
 - (A) Increasing breathlessness between the 12th and 24th week
 - (B) Tachycardia
 - (C) A fall in diastolic pressure
 - (D) Displacement of the apex beat to the left
- 120. An acute attack of migraine can be relieved or aborted by the early administration of:
 - (A) Methysergide
 - (B) Ergotamine tartrate
 - (C) Diazepam
 - (D) Reserpine
- 121. Upper to lower segment ratio of a normal newborn is:
 - (A)⊕ 1,3.:1.0 թանթան
 - (B) 1.5:1.0 as 26.648 6

- · (C) 1.7:1.0
- : -- // **(D)** 4.9:4.0 -- // -- //
 - 122. A child of 24 months can make tower of:
 - (A) 4 cubes
 - (B) 7 cubes
 - (C) 9 cubes
 - (D) 10 cubes
 - 123. A child copies a circle by:
 - (A) 2 years
 - (B) $2\frac{1}{2}$ years
 - (C) 3 years
 - (D) 4 years
 - 124. The fetal hemoglobin (HbF) is:
 - (A) Alpha2 Beta2
 - (B) Alpha2 Epsilon2
 - (C) Alpha2 Delta2
 - (D) Alpha2 Gamma2
- 125. Children with pica are at increased risk of intoxication with:
 - (A) Copper
 - (B) Iron
 - (C) Mercury
 - (D) Lead
- 126. Which among the following tissues of the body exhibit maximum rate of growth between 4 and 8 years of age?
 - (A) Brain
 - (B) Gonads
 - (C) Muscles
 - (D) Lymphoid tissue

WD – 1A/80

(15)





127.	In iron deficiency state, earliest to decrease is:		132.	Secondary naemorrnage occurs after:	
٠	(A)	Hemoglobin		(A) 1 day	
	(A) _. (B)	Hematocrit		(B) 3 days	
	(C)	Serum iron		(C) 5 days	
	(C) (D)	Serum ferritin		(D) 7 days	
128.	Myocarditis as a complication of diphtheria generally occurs by:		133.	The damage control surgery includes all except : (A) Arrest haemorrhage	
	(A)	2nd week		(B) Control Sepsis	
	(B)	3rd week		(C) Protect from further injury	
	(C)	4th week		(D) Definitive repair	
	(D)	5th week	134.	I. The following complications from	
129.	"Double Bubble" sign is the characteristic of:			massive blood transfusion may occur except:	
	(A)	Oesophageal atresia		(A) Hyperthermia	
	(B)	Duodenal atresia		(B) Hypothermia	
	(C)	Ileal atresia		(C) Hypokalaemia	
	•	Pyloric stenosis		(D) Hyperkalaemia	
130	(D) Defi	ciency of is	135.	5. All the statements about gas	
	responsible for neonatal seizure.			gangrene are correct except: (A) Immunocompromised patients	
	(A)	Vitamin B1		are at risk	
	(B)	Vitamin B12		(B) Smell is not characteristic	
	• •	Pyridoxine		(C) Gas is characteristic	1
	(D)	Vitamin E		(D) Caused by C-perfringens	
131.	com	following are the abdominal plications of Peritonitis except :	136.	5. The high risk lesion associate with malignant transformation in oral cavity is:	
. '	(A)	Residual abscess	- 	(A) Oral lichenplanus	
	(B)	Inflammatory mass		(B) Syphilitic glossitis	
* ,	(C)	Portal Pyaemia		(C) Oral Submucous fibrosis	
	(D)	Overactive bowel		(D) Erythroplakia	
WD:	- 1 A /	80 (16)	Contd.	





- 137. The features of Pelvic abscess are all except:
 - (A) Tachycardia
 - (B) Malaise
 - (C) Swinging Pyrexia
 - (D) Generalised abdominal tenderness
- 138. All are crystalloid solutions except:
 - (A) Ringer's lactate
 - (B) Hartmann's solution
 - (C) Albumen
 - (D) Normal saline
- 139. The Universal precautions include all except:
 - (A) Use of protective eye wear
 - (B) Use of boots
 - (C) Wearing double gloves
 - (D) All sharp instruments to be kept in one dish
- 140. The cause of Exudate Ascites is:
 - (A) Nephrotic Syndrome
 - (B) Chylous Ascites
 - (C) Cirrhosis
 - (D) Malnutrition
- 141. Ludwig's angina is:
 - (A) Inflammation of Submandibular region
 - (B) Tuberculous cervical lymphadenitis
 - (C) Actinomycosis of cervical lymph nodes
- (D) Reactive hyperplasia of cervical lymph nodes

- 142. The following statements about Mesenteric cyst are correct except:
 - (A) Cyst occurs most commonly in children
 - (B) It is twice as common in women as in men
 - (C) The most common presentation is of a painless abdominal swelling
 - (D) There is a zone of resonance around the cyst
- 143. Wolfe grafts are:
 - (A) Split thickness skin grafts
 - (B) Full thickness skin grafts
 - (C) Local flaps
 - (D) Distant flaps
- 144. Intravenous fluid resuscitation is required in children who sustained thermal burn injury with total body surface area of at least:
 - (A) 10%
 - (B) 20%
 - (C) 30%
 - (D) 40%
- 145. The treatment of Keloid includes all except:
 - (A) Pressure by local moulds
 - (B) Excision and steroid infection
 - (C) Laser
 - (D) Vitamin A

(17)





- 146. The Parkland formula for fluid therapy for 24 hours in burn patients is:
 - (A) Total percentage body surface area × wt(kg) × 2 = Volume (ml)
 - (B) Total percentage body surface area × wt(kg) × 4 = Volume (ml)
 - (C) Total percentage body surface area × wt(kg) × 6 = Volume (ml)
 - (D) Total percentage body surface area × wt(kg) × 8 = Volume (ml)
- 147. The complication of parotidectomy is:
 - (A) Sialadenosis
 - (B) Sarcoidosis
 - (C) Sialococle
 - (D) Sialadenitis
- 148. Meleney's Synergistic gangrene occurs in:
 - (A) Limb
 - (B) Scrotum
 - (C) Breast
 - (D) Abdominal wall
- 149. One of the component of systemic inflammatory response syndrome is:
 - (A) Hypothermia (< 36°C)
 - (B) Respiratory rate 16/min
 - (C) Pulse rate 80/min
 - (D) White cell count 5×109/L
- 150. Warthin's tumour is:
 - (A) Pleomorphic adenoma
 - (B) Lymphangioma

- (C) Adenolymphoma
- (D) Acinic cell carcinoma
- 151. All are the cardinal features of osteoarthritis except:
 - (A) Progressive cartilage destruction
 - (B) Sclerosis of surrounding bone
 - (C) Capsular fibrosis
 - (D) Intraarticular cyst formation
- 152. All are true about osteoarthrosis except:
 - (A) Gradual onset
 - (B) Reduced activity on scintigraphy
 - (C) Self limiting
 - (D) Focal changes on MRI
- 153. All are the factors adversely affecting the bone mass except:
 - (A) Early onset menopause
 - (B) Long term medication with antiepileptics
 - (C) Smoking
 - (D) Long term intake of bisphosphonates
- 154. One of the following is not a feature of Osteomalacia:
 - (A) Looser's zone
 - (B) Muscle weakness
 - (C) Decreased alkaline phosphatase
 - (D) Decreased serum phosphorus





- 155. It is not a feature of Renal Osteodystrophy:
 - (A) Epiphyseolysis
 - (B) Osteosclerosis in axial skeleton
 - (C) Rugger-Jersey spine
 - (D) Hypophosphataemia
- 156. One of the following is not a feature of Osteogenesis imperfecta:
 - (A) Fragility fracture
 - (B) Blue sclerae
 - (C) Crumbling teeth
 - (D) Osteosclerosis
- 157. Chondrocaleinosis is not seen in:
 - (A) Hyperparathyroidism
 - (B) Haemochromatosis
 - (C) Alkaptonuria
 - (D) Gout
- 158. Spontaneous osteonecrosis of knee is not characterized by:
 - (A) Involves elderly people
 - (B) Females are victim
 - (C) Osteoporosis
 - (D) Involvement of lateral femoral condyle
- 159. Intramembranous periosteal new bone formation occurs as a response to periosteal strepping due to:
 - (A) Tumour
 - (B) Trauma
 - (C) Infection
 - (D) Degenerative arthritis

- 160. Parathyroid Hormone:
 - (A) Secretion increases due to fall in serum calcium
 - (B) Directly increases absorption of calcium from intestine
 - (C) Controls hydroxylation of Vit. D metabolites in kidney
 - (D) Promotes osteoelastic bone resorption
- 161. X-Ray Paranasal sinus Caldwell view is best for visualization of which paranasal sinus?
 - (A) Maxillary sinus
 - (B) Frontal sinus
 - (C) Sphenoid sinus
 - (D) Ethmoid sinus
- 162. Which one of the following is not an elastic cartilage?
 - (A) Epiglottis
 - (B) Vocal process of arytenoids
 - (C) Apex of arytenoids
 - (D) Cuneiform cartilage
- 163. A 12 year old post op tonsillectomy patient comes to ENT clinic with fever and bleeding from tonsillectomy site at 10th post op day. The next line of management is:
 - (A) Immediately post for surgical exploration for bleeding site and cauterize
 - (B) Admit the patient and start on broad spectrum iv antibiotics
 - (C) External cold compression with ice pack, if uncontrolled then exploration
 - (D) Mouth gargle with hydrogen peroxide





- 164. Gradenigo's syndrome consists of all except:
 - (A) Retro-orbital pain
 - (B) Proptosis
 - (C) Strabismus
 - (D) Profuse otorrhoea
- 165. Superior laryngeal nerve supplies:
 - (A) Motor supply for Cricothyroid muscle
 - (B) Sensory supply for larynx below vocal cord
 - (C) Motor supply to thyroarytenoid
 - (D) Sensory supply to thyroid perichondrium
- 166. Which of the following is contraindicated in primary glaucoma?
 - (A) Atropine
 - (B) Pilocarpine
 - (C) Adrenaline
 - (D) Eserine
- 167. A newborn is invariably:
 - (A) Hypermetropic
 - (B) Myopic
 - (C) Astigmatic
 - (D) Aphakic
- 168. In retinoscopy using a plane mirror at 1 meter distance, when the mirror is tilted to the right the shadow in the pupil moves to the left in:
 - (A) Hypermetropia
 - (B) Myopia more than 1 D
 - (C) Emmetropia
 - (D) Myopia less than 1D

- 169. LASIK stands for:
 - (A) Laser augmented in situ keratomileusis
 - (B) Laser assisted in situ keratomileusis
 - (C) Light amplified in situ keratomileusis
 - (D) Light assisted in situ keratomileusis
- 170. Cherry red spot occurs in all EXCEPT:
 - (A) Central Retinal Vein Occlusion
 - (B) Central Retinal Artery
 Occlusion
 - (C) Tay-sach's disease
 - (D) Neimann-Pick disease
- 171. The diagnostic technique of choice for placental localization is:
 - (A) Ultrasonography
 - (B) MRI
 - (C) CT Scan
 - (D) Color Doppler Flow Study
- 172. The following method is used for diagnosing fetal anaemia in Rh incompatibility:
 - (A) Fetal MCA Doppler
 - (B) Fetal ductus venosus Doppler
 - (C) Uterine artery Doppler
 - (D) Umbilical artery Doppler
- 173. One of the following is NOT a complication of transverse life:
 - (A) Arm prolapse
 - (B) Cord prolapse
 - (C) Shoulder dystocia
 - (D) Obstructed labor

.





- 174. Risk of persistent Gestational Trophoblastic Disease (GTD) for complete mole is:
 - (A) 1% 5%
 - (B) 5% 10%
 - (C) 10% 15%
 - (D) 15% 20%
- 175. One of the following is **NOT** a predictor of successful VBAC:
 - (A) Fetal birth weight
 - (B) Amount of liquor
 - (C) Prior vaginal delivery
 - (D) Non recurring indication of previous caesarean section
- 176. The following drug is contraindicated during pregnancy:
 - (A) Labetalol
 - (B) Nifedipine
 - (C) Enalapril
 - (D) Hydralizine
- 177. One of the following is NOT a complication of preeclampsia:
 - (A) Acute kidney injury
 - (B) Abruptio placentae
 - (C) HELLP Syndrome
 - (D) Polyhydramnios
- 178. One of the following statements about atonic PPH is **WRONG**:
 - (A) Multiparity is a risk factor
 - (B) Active management of third stage of labor reduces the blood loss

- (C) Total hysterectomy with bilateral salpingo-oophorectomy is the treatment of choice
- (D) Uterine artery embolization is one of the methods for controlling PPH
- 179. Sinusoidal fetal heart rate is observed in the following condition:
 - (A) Post date pregnancy
 - (B) Preterm fetus
 - (C) Fetal anaemia
 - (D) Anencephalic fetus
- 180. Congenital anomalies are more common in the following conditions

 Except:
 - (A) Breech presentation
 - (B) Maternal diabetes
 - (C) Maternal hypertension
 - (D) Twins
- 181. The secondary mound is formed in the female breast in Tanner stage:
 - (A)
 - (B) II
 - (C) III
 - (D) IV
- 182. One of the following conditions is **NOT** responsible for ulcerative lesions on the vulva:
 - (A) Human papillomavirus infection
 - (B) Syphilis l
 - (C) Genital herpes
 - (D) Granuloma inguinale

41, 5





- 183. One of the following is **NOT** an indication for Intra Uterine Insemination (IUI):
 - (A) Oligospermia
 - (B) Unexplained infertility
 - (C) Premature ovarian failure
 - (D) Premature ejaculation
- 184. One of the following is a contraindication for methotrexate in ectopic pregnancy:
 - (A) Gestational sac 3 cm 5 cm
 - (B) Unruptured ectopic pregnancy
 - (C) Cervical pregnancy
 - (D) Heterotopic pregnancy
- 185. Serum FSH is increased in the following condition:
 - (A) Hyperprolactinemia
 - (B) Turner Syndrome
 - (C) Delayed menarchae
 - (D) Anorexia Nervosa
- 186. One of the following is NOT a sequelae of PCOS:
 - (A) Endometrial cancer
 - (B) Ovarian cancer
 - (C) Breast cancer
 - (D) Diabetes
- 187. The following statement about dermoid cyst of the ovary is FALSE:
 - (A) 5% 10% of all cystic tumors
 - (B) Frequently associated with mucinous cystadenoma

- (C) Common in young girls (< 20 years)
- (D) Epidermoid carcinoma can occur in < 2%
- 188. One of the following is **NOT** used for treating pre-invasive lesions of the cervix:
 - (A) Cryosurgery
 - (B) Hysterectomy
 - (C) Conization
 - (D) Chemoradiation
- 189. One of the following is NOT an indication for Colposcopy:
 - (A) High grade squamous intraepithelial lesion (HSIL)
 - (B) VIA positive
 - (C) Abnormal vulvar area
 - (D) Growth on the cervix
- 190. In endometrial carcinoma the serosa of the uterus is involved in FIGO stage:
 - (A) II-A
 - (B) · III−A
 - (C) I-B
 - (D) III-C
- 191. In a Respiratory cycle, at the end of Inspiration:
 - (A) The Intrapleural Pressure is most negative
 - (B) Alveolar Pressure is greater than Atmospheric Pressure
 - (C) The air flow reaches the Peak
 - (D) The Alveolar Pressure is around + 5 cm H₂O





- 192. Which of the following is FALSE regarding Atropine?
 - (A) May cause bradycardia
 - (B) Has a shorter duration of action than glycopyrolate
 - (C) Increases the Anatomical dead space
 - (D) Have both muscarinic and nicotinic effects
- 193. Which of the following is TRUE regarding Desflurane?
 - (A) Is 17 times more potent that Sevoflurane
 - (B) Is eliminated faster than Nitrous Oxide
 - (C) Is safe to use in patients with Malignant Hyperthermia
 - (D) Require a TEC 7 vaporiser for delivery of inhalational agent
- 194. Which of the following is NOT an inhalational anaesthetic delivery system?
 - (A) Aladin Cassette
 - (B) DIVA
 - (C) AnaConDa
 - (D) Lucas system
- 195. The 'CE' symbol noted in an endotracheal tube denotes:
 - (A) Conformittae Essentias
 - (B) Electromagnetic Compatible
 - (C) Conformite Europeene
 - (D) Certified Export

- 196. In Korsakoff Syndrome, pathological lesions are found in :
 - (A) Caudate nucleus
 - (B) Mamillary bodies
 - (C) Nucleus basalis of meynert
 - (D) Medial septal nucleus
- 197. The lesions of dorsolateral prefrontal cortex produce:
 - (A) Apathetic syndrome
 - (B) Euphoric syndrome
 - (C) Akinetic syndrome
 - (D) Parkinson's disease
- 198. A neuron utilizes the same neurotransmitter at all its terminals is called:
 - (A) Dale's principle
 - (B) Kindling effect
 - (C) Flooding
 - (D) Operant Conditioning
- 199. Negative views regarding onself, the world and the future is called:
 - (A) Cognitive triad
 - (B) Covert sensitization
 - (C) Implosion
 - (D) Mania
- 200. In phototherapy, the patient is exposed to a bright artificial light source daily for about:
 - (A) One hour before dawn
 - (B) Four hours before dawn
 - (C) Three hours before dawn
 - (D) Two hours before dawn





SPACE FOR ROUGH WORK



